

Chemical Examination of the Roots of *Cissampelos Pareira* Linn. IV.¹ Structure and Stereochemistry of Hayatin

A. K. BHATNAGAR, S. BHATTACHARJI, A. C. ROY, S. P. POPLI,
AND M. L. DHAR

Central Drug Research Institute, Lucknow, India


Received August 29, 1966

The isolation of hayatin [mp 298–303° dec, $[\alpha]_D 0^\circ$ (pyridine)] from the roots of *Cissampelos pareira* Linn. was described earlier by Bhattacharji, *et al.*^{2,3} The reported curariform activity^{4–7} of its dimethiodide led to an investigation of its chemical structure.

Sodium and liquid ammonia reduction of hayatin diethyl ether was undertaken with a view to establish, from the fragments obtained, the structure of hayatin as well as to determine the stereochemistry at the two asymmetric centers in the molecule.

Hayatin diethyl ether when subjected to sodium and liquid ammonia reduction gave a nonphenolic compound [mp 64°, $[\alpha]_D 0^\circ$ (CHCl₃), mass spectrum M^+ 355 (C₂₂H₂₉NO₃)] and a phenolic compound, [$[\alpha]_D 0^\circ$ (methanol)] which could not be crystallized. However, the R_f value of the phenol on tlc on silica gel (benzene–ethyl acetate–diethylamine, 7:2:1) was identical with that of N-methylcoclaurine. The O,O'-dimethyl methiodide of this compound, mp 131°, and the O,O'-diethyl methiodide derivative, mp 186°, were identical with samples of *dl*-O,O',N-trimethylcoclaurine methiodide and O,O'-diethyl N-methylcoclaurine methiodide, respectively, in respect to melting point, mixture melting point, and infrared spectra. This establishes the structure of the phenolic fragment as N-methylcoclaurine (I).

The structure of the nonphenolic fragment was determined as follows. Its mass spectrum showed abundant ions at m/e 220 (base peak), 204, and 190, a very weak molecular ion peak (M^+) at 355 (C₂₂H₂₉NO₃), and a weak peak at m/e 354 ($M - 1$). The base peak corresponded to the ion II or III and m/e 204 and 190



ions could presumably be formed by a further loss of a methyl or ethyl, respectively, with hydrogen transfer. That the m/e 220 ion was II and not III was proved by the nmr spectrum of the nonphenolic fragment which showed a singlet for three protons at τ 6.18 (6-OMe) and four protons in the region 5.93–6.32 (7,4'-(OCH₂-CH₃)₂). There were no signals at τ 6.4–6.5 which clearly showed the absence of an OMe at the 7 position of

the tetrahydroisoquinoline ring.⁸ The other features of the nmr spectrum were as follows: (i) six aromatic protons, four forming an A₂B₂ pattern with doublets centered at τ 2.95 and 3.22 ($J = 9$ cps) and two as singlets at 3.52 and 3.98, respectively; that the proton at τ 3.98 was the one at C-8 would be in consonance with the shielding by the bottom aromatic ring; (ii) two triplets of three protons each at τ 8.61 and 8.70, respectively [$J = 7.0$ cps, (OCH₂CH₃)₂]; and (iii) three protons for $>N \cdot Me$. All this data led to the formulation of the nonphenolic fragment as 7,4'-O,O-diethyl-N-methylcoclaurine (IV).

This was confirmed by establishing the identity of the methiodide of IV, mp 186°, with an authentic sample of *dl*-7,4'-O,O-diethyl-N-methylcoclaurine methiodide.

The confirmation of the phenolic and nonphenolic fragments as I and IV, respectively, and the fact that both hayatin and its sodium and liquid ammonia reduction products are optically inactive clearly establish the structure of hayatin as *dl*-bebeerine. This structure is fully supported by the nmr and mass spectra of hayatin and its diethyl ether.

It is pertinent to point out that the biological activity reported for hayatin dimethiodide agrees very closely with the computed values for *dl*-bebeerine dimethiodide on the basis of reported activities of the corresponding quaternary compounds of *d*- and *l*-bebeerine.⁹

Experimental Section

Hayatin Diethyl Ether.—Hayatin was ethylated with triethyl anilinium chloride in the usual manner, when a mixture of mono- and diethylated bases was obtained. The two were separated and purified by column chromatography on neutral alumina using benzene–chloroform mixture as the eluent. O,O'-Diethylhayatin was crystallized from methanol, mp 160° (yield 48%). *Anal.* Calcd for C₃₀H₄₆N₂O₆: C, 73.84; H, 7.07; N, 4.30. Found: C, 74.08; H, 7.12; N, 4.47.

Monoethylated hayatin was crystallized from methanol, mp 234° (yield 30%). *Anal.* Calcd for C₂₈H₄₂N₂O₆: C, 73.31; H, 6.75; N, 4.50. Found: C, 73.50; H, 6.90; N, 4.25.

Sodium and Liquid Ammonia Fission of O,O'-Diethylhayatin.—A solution of O,O'-diethylhayatin (500 mg), in a mixture of dry benzene (20 ml) and toluene (20 ml), was added dropwise to stirred liquid ammonia (500 ml) containing sodium (0.5 g). More sodium was added (total 1.5 g) until the blue color persisted. Stirring was continued for 2 hr and the mixture was then allowed to stand overnight and the ammonia was allowed to evaporate. The phenolic and nonphenolic fragments were then separated in the usual way. The latter, obtained in the form of fine plates, was recrystallized from methanol, yield 180 mg, mp 58° rising to 64° on drying at 40° under high vacuum; methiodide (methanol-ether) mp 186°. *Anal.* Calcd for C₂₂H₂₉NO₃: N, 2.81. Found: N, 2.68. Base oxalate (methanol-ether) had mp 188°. *Anal.* Calcd for C₂₄H₃₁NO₇·0.5H₂O: C, 63.43; H, 7.04; N, 3.08. Found: C, 63.15; H, 7.12; N, 3.24.

The phenolic portion (230 mg) was chromatographed on alumina (neutral). Elution with chloroform–methanol (49:1) gave a product (110 mg) which could not be crystallized. A part of this (70 mg) was methylated with diazomethane and then converted into the methiodide which was crystallized (methanol-ether), mp 131°, $[\alpha]_D (0^\circ)$ (methanol), yield 60 mg. A mixture melting point and infrared spectrum showed it to be identical with O,O',N-trimethylcoclaurine methiodide.

The rest of the phenolic fraction (40 mg) was ethylated with triethylanilinium chloride and the product after purification on alumina column was converted into the methiodide which was

(8) M. Tomita, T. Shingu, K. Fujitani, and H. Furukawa, *Chem. Pharm. Bull. (Tokyo)*, **13**, 921 (1965).

(9) R. B. Barlow, *Introduction to Chemical Pharmacology*, 2nd ed. Methuen and Co., Ltd., London, 1964, p 123.

(1) Part III. A. K. Bhatnagar and S. P. Popli, *Indian J. Chem.*, in press.
(2) S. Bhattacharji, V. N. Sharma, and M. L. Dhar, *J. Sci. Ind. Res. (India)*, **11B**, 81 (1952).

(3) S. Bhattacharji, V. N. Sharma, and M. L. Dhar, *ibid.*, **15B**, 363 (1956).

(4) S. N. Pradhan, C. Ray, and K. S. Varadan, *Current Sci. (India)*, **21**, 172 (1952).

(5) S. N. Pradhan and N. N. De, *Brit. J. Pharmacol.*, **8**, 399 (1953).

(6) S. N. Pradhan and N. N. De, *Arch. Intern. Pharmacodyn.*, **120**, 136 (1959).

(7) S. N. Pradhan, K. Pande, and R. P. Badola, *Brit. J. Anaesthesia*, **36**, 604 (1964).

Journal of Organic Chemistry 32, 819 (1967)

crystallized from methanol-ether (yield 20 mg), mp 186°. The methiodide was found to be identical with the methiodide of the nonphenolic fragment in respect to melting point, mixture melting point, and infrared spectrum.

Registry No.—Hayatin, 1381-51-7.

Acknowledgment.—Our thanks are due to Dr. T. R. Govindachari and Professor A. R. Battersby, F. R. S., for useful discussions and Dr. R. S. Kapil for determination of mass spectra.