

Concise synthesis of two trisaccharide analogs related to the glycone constituent of Phanoside, a novel insulin releasing natural product[†]

Geetanjali Agnihotri, Pintu Kumar Mandal and Anup Kumar Misra*

Medicinal and Process Chemistry Division, Central Drug Research Institute, Chattar Manzil Palace, Lucknow 226001, UP, India

Abstract— Two trisaccharides (**2** and **3**) related to the glycone part of Phanoside, an insulin release stimulator have been synthesized in excellent yield.

1. Introduction

Diabetes has been classified as insulin dependent (type 1) and non insulin dependent (type 2). Non insulin dependent diabetes (type 2) diabetes is the commonest form of diabetes constituting 90% of the diabetic population.^{1,2} Its prevalence in more affluent societies is spectacular and of general concern.³ It has broken the age barrier and also appears in younger people. Though its management through drugs is possible, drugs come together with serious side effects. Natural products can provide a better alternative as more effective anti-diabetics with less side effects. Recently, it has been reported that Phanoside⁴ (**1**, Figure 1), a natural product isolated from the plant *Gynostemma pentaphyllum* is effective in stimulating insulin release, and is much more potent than well known anti-diabetic drugs, such as sulphonylureas.⁴ As claimed in the original report, Phanoside contains a branched trisaccharide glycone consisting of D-glucose, D-rhamnose and D-lyxose.




Figure 1: Structure of Phanoside (**1**) and trisaccharide analogs of the glycone (**2** and **3**)

However, L-rhamnose is quite abundant in plant natural products and D-rhamnose is occasionally found in the microbial origin. Although the structural elucidation for the presence of D-rhamnose moiety instead of L-rhamnose was not clearly stated in the original paper, we were interested in synthesizing some analogous trisaccharide moieties related to the glycone part of Phanoside containing a L-rhamnose moiety. Oligosaccharides containing D-lyxose are also scarce in nature. In order to understand the role of the glycone part of Phanoside in enhancing the insulin release, it is essential to have higher quantities of several trisaccharide analogs related to the glycone part of Phanoside. Hence it was thought to develop a synthetic route for the synthesis of two trisaccharide analogs (**2** and **3**) related to the glycone part of Phanoside.

2. Results and discussion

The synthesis of target trisaccharide (**2**) as its 4-methoxyphenyl glycoside is demonstrated in Scheme 1. Treatment of D-lyxose tetraacetate (**4**) with 4-methoxy phenol in the presence of trifluoromethanesulfonic acid afforded 4-methoxy phenyl glycoside (**5**). Compound **5** was converted to compound **6** following a sequence of reactions consisting of deacetylation using sodium methoxide,⁵ isopropylideneation⁶ using 2,2-dimethoxypropane and *p*-toluenesulfonic acid and benzylation using benzyl bromide under phase transfer conditions.⁷ Removal of the isopropylidene group⁸ from compound **6** using 80% aq.

Keywords: Carbohydrate, glycosylations, natural products, insulin, phanoside.

[†] C.D.R.I communication no. 7049.

*Corresponding author. Tel.: +91-522-2612411-18, Ext. 4385; Fax: +91-522-2623405; e-mail: akmisra69@rediffmail.com

acetic acid furnished compound **7**. Initially, a regioselective glycosylation strategy⁹ was attempted to prepare disaccharide **8**, through the formation of stannylidene acetal of compound **7**, followed by treatment with peracetylated glucosyl bromide. After several trials, no disaccharide **8** could be isolated and starting material was recovered. Another regioselective glycosylation was carried out using ethyl 2,3,4,6-tetra-*O*-acetyl-1-thio- β -D-glucopyranoside under *N*-iodosuccinimide-triflic acid promoted glycosylation,¹⁰ but only some orthoester **9** could be isolated instead of disaccharide **8**. Following literature reports,¹¹ the rearrangement of orthoester **9** to the required disaccharide **8** also failed and compound **7** was recovered. It was then decided to selectively protect the 3-hydroxy group of compound **7** with the 4-methoxybenzyl group via stannylidene acetal formation. Unfortunately, a mixture of products was isolated upon treatment of compound **7** with dibutyltin oxide followed by treatment with 4-methoxybenzyl chloride. Finally, stannylidene acetal formation of compound **7** followed by treatment with benzoyl chloride¹² yielded selectively benzoyl protected D-lyxose derivative **10** in excellent yield. Glycosylation of compound **10** with ethyl 2,3,4-tri-*O*-benzyl-1-thio- α -L-rhamnopyranoside (**11**), prepared from L-rhamnose in three steps, with *N*-iodosuccinimide and triflic acid¹⁰ afforded the disaccharide derivative **12**. The presence of two anomeric signals at 97.4 (C-1), 97.0 (C-1') in the ¹³C spectrum confirmed the formation of **12**. Following an earlier literature report,¹³ the configuration of the L-rhamnosyl linkage of compound **12** was confirmed by the partially decoupled ¹³C NMR spectrum, in which the C-1/H-1 coupling constant ($J_{C-1/H-1}$) for L-rhamnosyl linkage was found to be 172.0 Hz indicating the formation of a α -linkage. Removal of the benzoyl group using sodium methoxide gave compound **13**, which on glycosylation with phenyl 2,3,4,6-tetra-*O*-benzoyl- α -D-glucopyranoside (**14**)¹⁴ using *N*-iodosuccinimide and trifluoromethane sulfonic acid furnished trisaccharide derivative **15**. The presence of three anomeric signals at 101.8 (C-1''), 99.7 (C-1), 97.4 (C-1') in the ¹³C spectrum, confirmed the formation of trisaccharide derivative **15**. Conventional debenzoylation followed by hydrogenolysis¹⁵ of trisaccharide **15** afforded target trisaccharide **2** in 62% yield.

In a separate experiment, another analogous trisaccharide **3** as its 4-methoxyphenyl glycoside was prepared following Scheme 2. Glycosylation of compound **7** with ethyl 2,3,4-tri-*O*-acetyl-1-thio- α -L-rhamnopyranoside (**16**)¹⁶ using *N*-iodosuccinimide and trifluoromethane sulfonic acid furnished analogous trisaccharide derivative **17**. Formation of compound **17** was confirmed by ¹H, ¹³C and partially decoupled ¹³C NMR spectra. Saponification followed by hydrogenolysis of trisaccharide **17** furnished the trisaccharide **3**, which was confirmed by NMR and mass spectroscopy.

3. Conclusions

In summary, two trisaccharide analogs related to the glycone constituent of Phanoside, a novel insulin releasing natural product have been synthesized as 4-methoxyphenyl glycosides in a very concise manner.

4. Experimental

General Procedure

All the reactions were monitored by thin layer chromatography over silica gel coated TLC plates. The spots on TLC were visualized by warming ceric sulphate (2% Ce(SO₄)₂ in 2M H₂SO₄) sprayed plates on a hot plate. Silica gel 230-400 mesh was used for column chromatography. ¹H and ¹³C NMR were recorded on a Bruker Advance DPX 300 MHz using TMS as internal reference. Chemical shift values were expressed in δ ppm. Elemental analysis was carried out on a Carlo ERBA-1108 analyzer. Optical rotations were measured at 25°C on a Rudolf Autopol III polarimeter. Commercially available grades of organic solvents of adequate purity are used in many reactions.

4-Methoxyphenyl 2,3,4-tri-*O*-acetyl- α -D-lyxopyranoside (5): To a solution of 1,2,3,4-tetra-*O*-acetyl- α -D-lyxopyranose (**4**, 6.0 g, 18.8 mmol) and 4-methoxy phenol (2.81 g, 22.6 mmol) in dry CH₂Cl₂ (25 mL) was added trifluoromethane sulfonic acid (150 μ L) at 0 °C and the reaction mixture was allowed to stir at 10 °C for 2 h. The reaction mixture was diluted with CH₂Cl₂ and the organic layer was washed successively with water, aq. NaHCO₃ and water, dried (Na₂SO₄), and concentrated under reduced pressure. Purification of the crude reaction product by column chromatography on silica gel using hexane–EtOAc (4:1) furnished pure compound **5** (6.5 g, 90%) as thick syrup; $[\alpha]_D^{25} + 31.1$ (*c* 1.2, CHCl₃); IR (neat): 1715, 1509, 1372, 1222, 1044, 831 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.00-6.96 (m, 2 H, ArH), 6.83-6.73 (m, 2 H, ArH), 5.50 (dd, *J* = 9.9 and 3.6 Hz, 1 H, 3-H), 5.39 (dd, *J* = 3.3 and 2.4 Hz each, 1 H, 2-H), 5.30 (d, *J* = 2.4 Hz, 1 H, 1-H), 5.27-5.19 (m, 1 H, 4-H), 3.90 (dd, *J* = 11.1 and 5.7 Hz, 1 H, 5-H_a), 3.77 (s, 3 H, OCH₃), 3.76-3.71 (m, 1 H, 5-H_b), 2.18, 2.04 (2s, 9 H, 3 COCH₃); ¹³C NMR (CDCl₃, 75 Hz): δ 169.5, 169.4, 169.3 (3 COCH₃), 155.4, 149.9, 117.9 (2 C), 114.6 (2 C), 96.8 (C-1), 69.5, 68.5, 66.8, 60.2 (C-5), 55.4 (OCH₃), 20.7 (3 C, COCH₃); ESI-MS: *m/z* = 405.1 [M+Na]⁺; Anal. Calcd. for C₁₈H₂₂O₉ (382.1): C, 56.54; H, 5.80; Found: C, 56.30; H, 6.10.

4-Methoxyphenyl 4-*O*-benzyl-2,3-*O*-isopropylidene- α -D-lyxopyranoside (6): A solution of compound **5** (5.0 g, 13.1 mmol) in 0.05 M sodium methoxide in methanol (30.0 mL) was stirred at room temperature for 3 h. The reaction mixture was neutralized with Dowex 50W-X8 (H⁺), filtered and the filtrate was evaporated to dryness to give a glassy solid in quantitative yield (3.16 g). The dried mass thus obtained was dissolved in anhydrous DMF (15 mL) and 2,2-dimethoxy propane (2.4 mL,

18.5 mmol) was added to it followed by addition of *p*-toluene sulphonic acid (~100 mg) to make solution acidic (pH~2). After stirring at room temperature for 6 h the reaction mixture was neutralized with triethylamine and filtered through the celite bed. The filtrate was evaporated to dryness and the crude mass was purified over SiO₂ using hexane-EtOAc (3:1) to afford pure 4-methoxyphenyl 2,3-*O*-isopropylidene- α -D-lyxopyranoside (2.77 g, 71%), which was dissolved in CH₂Cl₂ (20 mL) and benzyl bromide (1.6 mL, 14.0 mmol) followed by 50% aq. KOH solution (5 mL) and tetrabutylammonium iodide (~100 mg) was added to it. After stirring at room temperature for 3 h the reaction mixture was diluted with CH₂Cl₂ (50 mL). The organic layer was washed successively with water, aq. NaCl and water, dried (Na₂SO₄), and concentrated under reduced pressure. Column chromatography of the crude product over silica gel using hexane-EtOAc (5:1) furnished pure compound **6** (3.25 g, 90%) as syrup; [α]_D²⁵ + 60.0 (*c* 1.2, CHCl₃); IR (neat): 2359, 1597, 1375, 1225, 1073 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.33-7.20 (m, 5 H, ArH), 6.95-6.89 (m, 2 H, ArH), 6.79-6.73 (m, 2 H, ArH), 5.42 (d, *J* = 1.5 Hz, 1 H, 1-H), 4.72 (d, *J* = 12.0 Hz, 1 H, CH₂C₆H₅), 4.61 (d, *J* = 12.0 Hz, 1 H, CH₂C₆H₅), 4.34-4.30 (m, 1 H, 2-H), 4.27 (dd, *J* = 5.7 and 1.8 Hz, 1 H, 3-H), 3.73 (s, 3 H, OCH₃), 3.63-3.60 (m, 3 H, 4-H and 5-H_{ab}), 1.45, 1.37 (2 s, 6 H, C(CH₃)₂); ¹³C NMR (CDCl₃, 75 Hz): δ 155.1, 150.3, 138.2, 128.3 (2 C), 127.8 (2 C), 127.7, 117.8 (2 C), 114.6 (2 C), 109.3 (C(CH₃)₂), 97.0 (C-1), 77.5, 75.4 (PhCH₂), 74.5, 71.9, 59.5 (C-5), 55.4 (OCH₃), 28.1 (C(CH₃)₂), 26.5 (C(CH₃)₂); ESI-MS: *m/z* = 409.2 [M+Na]⁺; Anal. Calcd. for C₂₂H₂₆O₆ (386.2): C, 68.38; H, 6.78; Found: C, 68.20; H, 6.95.

4-Methoxyphenyl 4-*O*-benzyl- α -D-lyxopyranoside (7): Compound **6** (3.0 g, 7.77 mmol) was dissolved in 80% aq. acetic acid (25 mL) and the reaction mixture was stirred at 60 °C for 45 min. The reaction mixture was evaporated and co-evaporated with toluene under reduced pressure. Purification of the crude reaction product using a short pad of SiO₂ using hexane-EtOAc (3:1) as the eluant furnished compound **7** (2.5 g, 93%) as yellow syrup; [α]_D²⁵ + 50.2 (*c* 1.2, CHCl₃); IR (neat): 2927, 2368, 2595, 1352, 1029 673 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 7.31-7.25 (m, 5 H, ArH), 6.93 (d, *J* = 9.0 Hz, 2 H, ArH), 6.77 (d, *J* = 9.0 Hz, 2 H, ArH), 5.30 (br s, 1 H, 1-H), 4.64-4.60 (m, 2 H, CH₂Ph), 4.09-4.07 (m, 2 H, 2-H and 3-H), 3.75 (s, 3 H, OCH₃), 3.80-3.62 (m, 3 H, 4-H and 5-H_{ab}); ¹³C NMR (CDCl₃, 75 Hz): δ 155.1, 150.4, 137.9, 128.6 (2 C), 127.8 (3 C), 117.8 (2 C), 114.6 (2 C), 98.6 (C-1), 75.3, 72.5 (PhCH₂), 70.5, 70.3, 61.0 (C-5), 55.5 (OCH₃); ESI-MS: *m/z* = 369.1 [M+Na]⁺; Anal. Calcd. for C₁₉H₂₂O₆ (346.1): C, 65.88; H, 6.40; Found: C, 65.70; H, 6.62.

4-Methoxyphenyl 3-*O*-benzoyl-4-*O*-benzyl- α -D-lyxopyranoside (10): A solution of compound **7** (2.5 g, 7.22 mmol) and dibutyltin oxide (2.16 g, 8.67 mmol) in anhydrous methanol (20 mL) was refluxed for 3 h. The reaction mixture was concentrated and co evaporated with toluene several times. The dried mass was redissolved in dry toluene and benzoyl chloride (930 μ L, 7.92 mmol) was added to it and the reaction mixture was allowed to stir at room temperature for 8 h. The reaction was quenched by addition of methanol and reaction mixture was concentrated. Column chromatography of the crude product over silica gel using hexane-EtOAc (4:1) furnished pure compound **10** (2.6 g, 80%) as glassy solid; [α]_D²⁵ + 40.8 (*c* 1.2, CHCl₃); IR (KBr): 2938, 2365, 1695, 1594, 1286, 1093, 1006, 705 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 8.08-8.05 (m, 2 H, ArH), 7.61-7.56 (m, 1 H, ArH), 7.48-7.43 (m, 2 H, ArH), 7.28-7.14 (m, 5 H, ArH), 7.02 (d, *J* = 8.7 Hz, 2 H, ArH), 6.82 (d, *J* = 8.4 Hz, 2 H, ArH), 5.64 (dd, *J* = 8.1 and 3.0 Hz, 1 H, 3-H), 5.34 (d, *J* = 3.0 Hz, 1 H, 1-H), 4.65 (s, 2 H, CH₂Ph), 4.34 (t, *J* = 2.7, 2.7 Hz, 1 H, 2-H), 4.09-4.02 (ddd, *J* = 8.1, 5.4 and 2.7 Hz, 1 H, 4-H), 3.90-3.82 (m, 2 H, 5-H_{ab}), 3.78 (s, 3 H, OCH₃); ¹³C NMR (CDCl₃, 75 Hz): δ 165.4 (COPh), 155.1, 150.2, 137.9, 133.2 (2 C), 129.9, 129.8 (2 C), 128.4 (2 C), 128.3 (2 C), 127.8, 127.7, 117.9 (2 C), 114.5 (2 C), 98.9 (C-1), 73.3, 72.6, 72.3, 69.0, 61.8 (C-5), 55.4 (OCH₃); ESI-MS: *m/z* = 473.2 [M+Na]⁺; Anal. Calcd. for C₂₆H₂₆O₇ (450.1): C, 69.32; H, 5.82; Found: C, 69.14; H, 6.04.

4-Methoxyphenyl 2,3,4-tri-*O*-benzyl- α -L-rhamnopyranosyl-(1 \rightarrow 2)-3-*O*-benzoyl-4-*O*-benzyl- α -D-lyxopyranoside (12) : To a solution of compound **10** (1.0 g, 2.22 mmol) and compound **11** (1.3 g, 2.66 mmol) in anhydrous CH₂Cl₂ (15 mL) was added powdered MS 4Å (2.0 g) and the mixture was stirred under argon for 1 h. *N*-Iodosuccinimide (658 mg, 2.92 mmol) was added to the reaction mixture and cooled to -30°C. Trifluoromethanesulfonic acid (25 μ L, 0.29 mmol) was added to the reaction at -30 °C and the reaction mixture was allowed to stir for 30 min at -30 °C. The reaction mixture was filtered through a celite bed and the organic layer was washed with 5% aq. Na₂S₂O₃, satd. NaHCO₃, water in succession, dried (Na₂SO₄) and concentrated to a syrupy product. Column chromatography of the crude product over SiO₂ using hexane-EtOAc (7:1) afforded pure disaccharide derivative **12** (1.5 g, 78%) as syrup; [α]_D²⁵ -27.1 (*c* 1.2, CHCl₃); IR (neat): 2367, 1595, 1351, 1057 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz): δ 8.06-8.03 (m, 2 H, ArH), 7.54-7.50 (m, 1 H, ArH), 7.35-7.25 (m, 2 H, ArH), 7.02-6.96 (m, 2 H, ArH), 6.78 (d, *J* = 9.0 Hz, 2 H, ArH), 5.62 (dd, *J* = 8.4 and 3.0 Hz, 1 H, 3-H), 4.96 (br s, 1 H, 1-H), 4.90 (d, *J* = 11.1 Hz, 1 H, CH₂Ph), 4.79 (d, *J* = 12.3 Hz, 1 H, CH₂Ph), 4.69-4.60 (m, 5 H, 2 CH₂Ph, 1'-H), 4.58 (d, 12.6 Hz, 1 H, CH₂Ph), 4.53 (d, *J* = 11.0 Hz, 1 H, CH₂Ph), 4.33-4.28 (m, 1 H, 2-H), 4.0-3.92 (m, 1 H, 5'-H), 3.88 (s, 3 H, OCH₃), 3.86-3.84 (m, 2 H, 5-H_{ab}), 3.83 (t, *J* = 4.5, 4.5 Hz, 1 H, 2'-H), 3.81-3.78 (m, 1 H, 4-H), 3.76 (dd, *J* = 9.0, 3.1 Hz, 1 H, 3'-H), 3.51 (t, *J* = 9.3, 9.3 Hz, 1 H, 4'-H), 0.98 (d, *J* = 6.3 Hz, 3 H, CH₃); ¹³C NMR (CDCl₃, 75 Hz): δ 165.5 (COPh), 154.4, 151.1, 139.3-114.9 (Ar C), 97.4 (C-1), 97.0 (C-1'), 80.6, 80.0, 75.7, 75.3 (CH₂Ph), 73.7 (CH₂Ph), 73.2 (CH₂Ph), 72.8 (CH₂Ph), 72.7, 72.3 (2 C), 68.9, 62.3 (C-5), 57.0 (OCH₃), 17.8 (CH₃); ESI-MS: *m/z* 889.5 [M+Na]⁺; Anal. Calcd. for C₅₃H₅₄O₁₁ (866.3): C, 73.42; H, 6.28; found: C, 73.25; H, 6.50.

4-Methoxyphenyl 2,3,4-tri-*O*-benzyl- α -L-rhamno-pyranosyl-(1 \rightarrow 2)-4-*O*-benzyl- α -D-lyxopyranoside (13): A solution of compound **12** (1.5 g, 1.73 mmol) in 0.1 M sodium methoxide in methanol (10.0 mL) was stirred at room temperature for 3 h.

The reaction mixture was neutralized with Dowex 50W-X8 (H^+), filtered and the filtrate was evaporated to dryness to give disaccharide derivative **13** as syrup (1.31 g); $[\alpha]_D^{25} -16.0$ (c 1.2, $CHCl_3$); IR (neat): 2926, 2369, 1592, 1351 cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 7.32-7.18 (m, 20 H, ArH), 6.89 (d, $J = 9.0$ Hz, 2 H, ArH), 6.76 (d, $J = 9.0$ Hz, 2 H, ArH), 5.05 (d, $J = 3.8$ Hz, 1 H, 3-H), 4.93 (br s, 1 H, 1-H), 4.90 (d, $J = 11.1$ Hz, 1 H, CH_2Ph), 4.74-4.59 (m, 8 H, 4 CH_2Ph , 1'-H), 4.08-4.01 (m, 1 H, 5'-H), 4.0-3.96 (m, 1 H, 2-H), 3.86-3.80 (m, 2 H, 2'-H and 4-H), 3.76 (s, 3 H, OCH_3), 3.75 (dd, $J = 9.0, 2.7$ Hz, 1 H, 3'-H), 3.68-3.56 (m, 3 H, 5- $H_{a,b}$ and 4'-H), 1.32 (d, $J = 6.0$ Hz, 3 H, CH_3); ^{13}C NMR ($CDCl_3$, 75 Hz): δ 155.2-127.7 (Ar C), 117.9 (2 C), 114.5 (2 C), 98.5 (C-1), 97.8 (C-1'), 80.3, 79.4, 76.1, 75.4 (2 C, CH_2Ph), 75.2 (CH_2Ph), 72.9 (CH_2Ph), 72.4, 72.3, 70.2, 69.1, 61.8 (C-5), 55.4 (OCH_3), 18.1 (CH_3); ESI-MS: m/z 785.3 $[M+Na]^+$; Anal. Calcd. for $C_{46}H_{50}O_{10}$ (762.3): C, 72.42; H, 6.61; found: C, 72.25; H, 6.80.

4-Methoxyphenyl [2,3,4-tri-O-benzyl- α -L-rhamno-pyranosyl-(1 \rightarrow 2)]-[2,3,4,6-tetra-O-benzoyl- β -D-gluco-pyranosyl-(1 \rightarrow 3)]-4-O-benzyl- α -D-lyxopyranoside (15): To a solution of compound **13** (1.0 g, 1.31 mmol) and phenyl 2,3,4,6-tetra-O-benzoyl-1-thio- β -D-glucopyranoside (**14**; 1.1 g, 1.6 mmol) in anhydrous CH_2Cl_2 (10 mL) was added powdered MS 4 \AA (1.5 g) and the mixture was stirred under argon for 1 h. *N*-Iodosuccinimide (390 mg, 1.73 mmol) was added to the reaction mixture and cooled to $-30^\circ C$. Trifluoromethane sulfonic acid (15 μL , 0.17 mmol) was added to the reaction mixture at $-30^\circ C$ and the reaction mixture was allowed to stir for 30 min at $-30^\circ C$. After completion of the reaction, the reaction mixture was filtered through a celite bed and the organic layer was washed with 5% aq. $Na_2S_2O_3$, satd. $NaHCO_3$, water, dried (Na_2SO_4) and concentrated to a syrupy mass. Column chromatography of the crude product over SiO_2 using hexane-EtOAc (5:1) afforded pure trisaccharide derivative **15** (1.26 g, 72%); $[\alpha]_D^{25} + 10.6$ (c 1.2, $CHCl_3$); IR (neat): 2924, 2368, 1595, 1351, 1097, 708 cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 8.10-8.04 (m, 2 H, ArH), 7.85-7.80 (m, 4 H, ArH), 7.74-7.71 (m, 2 H, ArH), 7.52-7.46 (m, 4 H, ArH), 7.42-7.22 (m, 22 H, ArH), 7.20-7.07 (m, 4 H, ArH), 6.91-6.86 (m, 2 H, ArH), 6.57 (d, $J = 9.0$ Hz, 2 H, ArH), 6.46 (d, $J = 8.7$ Hz, 2 H, ArH), 5.88 (t, $J = 9.6, 9.6$ Hz, 1 H, 3''-H), 5.62 (t, $J = 9.6, 9.6$ Hz, 1 H, 2''-H), 5.42 (t, $J = 9.6, 9.6$ Hz, 1 H, 4''-H), 5.34 (d, $J = 8.1$ Hz, 1 H, 1''-H), 5.25 (br s, 1 H, 1-H), 5.05 (d, $J = 11.1$ Hz, 1 H, CH_2Ph), 4.82-4.70 (m, 4 H, 2 CH_2Ph , 3-H), 4.66 (br s, 1 H, 1'-H), 4.61 (d, $J = 11.6$ Hz, 1 H, CH_2Ph), 4.56-4.51 (m, 3 H, CH_2Ph), 4.39 (t, $J = 3.6, 3.6$ Hz, 1 H, 2-H), 4.30 (dd, $J = 12.0, 5.1$ Hz, 1 H, 6''- H_a), 4.17-4.10 (m, 2 H, 6''- H_b and 5'-H), 3.92 (dd, $J = 9.0, 2.7$ Hz, 1 H, 3'-H), 3.84-3.82 (m, 2 H, 5''-H and 2'-H), 3.81-3.77 (m, 1 H, 4-H), 3.76 (s, 3 H, OCH_3), 3.75-3.73 (m, 2 H, 5- $H_{a,b}$), 3.72-3.68 (m, 1 H, 4'-H), 1.42 (d, $J = 6.0$ Hz, 3 H, CH_3); ^{13}C NMR ($CDCl_3$, 75 Hz): δ 165.7, 165.6, 164.9, 164.8 (4 $COPh$), 138.8-114.3 (Ar C), 101.0 (C-1''), 99.5 (C-1), 98.7 (C-1'), 80.9, 80.5, 75.7, 75.4, 74.7, 73.4, 72.7 (2 C), 72.6 (2 C, CH_2Ph), 72.5, 71.2 (2 C, CH_2Ph), 69.4 (2 C), 63.2 (C-6''), 62.6 (C-5), 55.4 (OCH_3), 18.5 (CH_3); ESI-MS: m/z 1363.5 $[M+Na]^+$; Anal. Calcd. for $C_{80}H_{76}O_{19}$ (1340.5): C, 71.63; H, 5.71; found: C, 71.45; H, 5.98.

4-Methoxyphenyl [α -L-rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 3)]- α -D-lyxopyranoside (2): A solution of compound **15** (1.0 g, 0.75 mmol) in 0.1 M sodium methoxide in methanol (25.0 mL) was stirred at room temperature for 12 h. The reaction mixture was neutralized with Dowex 50W-X8 (H^+), filtered and the filtrate was evaporated to dryness to give a syrup, which was purified through Sephadex LH-20 using toluene as eluant to give 4-methoxyphenyl [2,3,4-tri-O-benzyl- α -L-rhamnopyranosyl-(1 \rightarrow 2)]- β -D-glucopyranosyl-(1 \rightarrow 3)]-4-O-benzyl- α -D-lyxopyranoside, which was dissolved in methanol (5 mL) and 20% $Pd(OH)_2-C$ (300 mg) was added to it. The reaction mixture was stirred under hydrogen at room temperature for 18 h. The mixture was filtered through a Celite bed and concentrated to give target trisaccharide **2** (260 mg; 61%) as syrup; $[\alpha]_D^{25} + 22$ (c 1.2, CH_3OH); 1H NMR (CD_3OD-D_2O , 300 MHz): δ 6.92-6.89 (m, 2 H, ArH), 6.77-6.73 (m, 2 H, ArH), 5.28 (d, $J = 4.8$ Hz, 1 H, 1'-H), 4.96 (br s, 1 H, 1-H), 4.47 (d, $J = 7.8$ Hz, 1 H, 1''-H), 4.10-4.03 (m, 2 H, 2'-H and 3'-H), 3.88-3.86 (m, 1 H, 4'-H), 3.84-3.83 (m, 1 H, 2-H), 3.82-3.77 (m, 3 H, 6''- $H_{a,b}$ and 4''-H), 3.64 (s, 3 H, OCH_3), 3.59-3.53 (m, 2 H, 3-H and 5'-H), 3.35-3.27 (m, 2 H, 3''-H and 4-H), 3.25-3.20 (m, 3 H, 5''-H and 5- $H_{a,b}$), 3.17-3.14 (m, 1 H, 2''-H), 1.22 (d, $J = 6.3$ Hz, 3 H, CH_3); ^{13}C NMR (CD_3OD-D_2O , 75 Hz): δ 154.0, 149.6, 116.4 (2 C), 112.9 (2 C), 101.8 (C-1''), 99.7 (C-1), 97.4 (C-1'), 77.1 (2 C), 75.4 (2 C), 75.0, 72.4 (2 C), 71.2, 69.7, 69.4, 67.9, 60.0 (2 C, C-5 and C-6''), 53.4 (OCH_3), 15.3 (CH_3); ESI-MS: m/z 587.2 $[M+Na]^+$; Anal. Calcd. for $C_{24}H_{36}O_{15}$ (564.2): C, 51.06; H, 6.43; found: C, 50.85; H, 6.65.

4-Methoxyphenyl [2,3,4-tri-O-acetyl- α -L-rhamno-pyranosyl-(1 \rightarrow 2)]-[2,3,4-tri-O-acetyl- α -L-rhamno-pyranosyl-(1 \rightarrow 3)]-4-O-benzyl- α -D-lyxopyranoside (17): To a solution of compound **7** (500 mg, 1.44 mmol) and ethyl 2,3,4-tri-O-acetyl- α -L-rhamnopyranoside (**16**; 1.2 g, 3.60 mmol) in anhydrous CH_2Cl_2 (10 mL) was added powdered MS 4 \AA (1.0 g) and the mixture was stirred under argon for 1 h. *N*-Iodosuccinimide (890 mg, 3.97 mmol) was added to the reaction mixture and it was cooled to $-30^\circ C$. Trifluoromethanesulfonic acid (35 μL , 0.39 mmol) was added and the reaction mixture was allowed to stir for 30 min at $-30^\circ C$. After completion of the reaction, the reaction mixture was filtered through a Celite bed and the filtrate was washed with 5% aq. $Na_2S_2O_3$, satd. $NaHCO_3$, water, dried (Na_2SO_4) and concentrated to a syrupy product. Column chromatography of the crude product over SiO_2 using hexane-EtOAc (3:1) afforded pure trisaccharide derivative **17** (872 mg, 68%); $[\alpha]_D^{25} - 26.1$ (c 1.2, $CHCl_3$); IR (neat): 2935, 2366, 1752, 1594, 1375, 1225, 1047 cm^{-1} ; 1H NMR ($CDCl_3$, 300 MHz): δ 7.28-7.14 (m, 5 H, ArH), 6.97 (d, $J = 8.7$ Hz, 2 H, ArH), 6.78 (d, $J = 9.0$ Hz, 2 H, ArH), 5.35-5.24 (m, 5 H, 1-H, 1'-H, 2'-H, 3'-H, 4'-H), 5.13-4.97 (m, 6 H, 1''-H, 2''-H, 3''-H, 4''-H, CH_2Ph), 4.20-4.09 (m, 4 H, 5'-H, 5''-H, 2-H, 3-H), 4.04-3.87 (m, 2 H, 5- $H_{a,b}$), 3.77 (s, 3 H, OCH_3), 3.67-3.60 (m, 1 H, 4-H), 2.18, 2.17, 2.16, 2.07, 2.06, 2.05 (6 s, 18 H, 6 $COCH_3$),

1.22-1.24 (m, 6 H, 2 CH_3); ^{13}C NMR (CDCl_3 , 75 Hz): δ 169.9, 169.7, 169.5, 169.4, 169.3, 169.2 (6 $\text{C}=\text{O}$), 155.1-114.4 (Ar C), 98.5 (C-1), 97.3 (C-1' and C-1''), 71.2, 70.6, 70.5 (2 C), 70.2 (C_6H_5), 69.8, 69.5, 69.1, 68.8, 68.7, 68.5, 67.1, 66.6 (C-5), 55.3 (OCH_3), 20.6 (2 C, 2 COCH_3), 20.5 (2 C, 2 COCH_3), 20.3 (2 C, 2 COCH_3), 17.3, 17.0 (2 CH_3); ESI-MS: m/z 913.3 $[\text{M}+\text{Na}]^+$; Anal. Calcd. for $\text{C}_{43}\text{H}_{54}\text{O}_{20}$ (890.3): C, 57.97; H, 6.11; found: C, 57.75; H, 6.30.


4-Methoxyphenyl [α -L-rhamnopyranosyl-(1 \rightarrow 2)]- [α -L-rhamnopyranosyl-(1 \rightarrow 3)]- α -D-lyxopyranoside (3): To a solution of compound **17** (800 mg, 0.9 mmol) in methanol, was added 20% $\text{Pd}(\text{OH})_2\text{-C}$ (100 mg) and reaction mixture was stirred under hydrogen at room temperature for 10 h. The mixture was filtered through celite bed and concentrated to give yellow syrup, which was dissolved in 0.1 M sodium methoxide in methanol (15.0 mL) and the reaction mixture was stirred at room temperature for 12 h. The reaction mixture was neutralized with Dowex 50W-X8 (H^+), filtered and the filtrate was evaporated to dryness to give a syrup, which was purified through Sephadex LH-20 using 80% aqueous EtOH as eluent to give pure trisaccharide **3** (320 mg, 65%) as a viscous liquid; $[\alpha]_D^{25} - 10$ (c 1.2, CH_3OH); ^1H NMR ($\text{CD}_3\text{OD-D}_2\text{O}$, 300 MHz): δ 6.92 (d, $J = 8.7$ Hz, 2 H, ArH), 6.76 (d, $J = 9.0$ Hz, 2 H, ArH), 5.15 (bs, 1 H, 1-H), 5.04 (d, $J = 2.6$ Hz, 2 H, 1'-H and 1''-H), 4.09-4.08 (m, 1 H, 2'-H), 3.99-3.98 (m, 1 H, 2''-H), 3.86-3.74 (m, 3 H, 3'-H, 3''-H and 2-H), 3.73-3.55 (m, 4 H, 4'-H, 4''-H, 3-H and 4-H), 3.65 (s, 3 H, OCH_3), 3.39-3.29 (m, 4 H, 5'-H, 5''-H and 5- $\text{H}_{a,b}$), 1.23-1.13 (m, 6 H, 2 CH_3); ^{13}C NMR ($\text{CD}_3\text{OD-D}_2\text{O}$, 75 MHz): 156.7, 149.1, 119.1 (2 C), 115.6 (2 C), 100.1 (2 C, C-1' and C-1''), 94.5 (C-1), 73.8 (2 C), 73.7 (2 C), 72.2, 72.1 (2 C), 72.0 (2 C), 70.5, 70.4, 68.8 (C-5), 56.1 (OCH_3), 18.1, 17.9 (2 CH_3); ESI-MS: m/z 571.2 $[\text{M}+\text{Na}]^+$; Anal. Calcd. for $\text{C}_{24}\text{H}_{36}\text{O}_{14}$ (548.2): C, 52.55; H, 6.62; found: C, 52.38; H, 6.85.

Acknowledgements


Instrumentation facilities from SAIF, CDRI are gratefully acknowledged. G.A. and P.K.M. thank CSIR, New Delhi for providing a Senior and Junior Research Fellowship respectively. This project was partly funded by Department of Science and Technology (DST), New Delhi (Project no. SR/FTP/CSA-10/2002), India.

References

- (1) Harris, M. Genetic environmental interaction in Diabetes mellitus. Eds. Mellish, J. S.; Hanna, J. B. *Experta Medica, Amsterdam*, **1982**, 70-76.
- (2) Zimmer, P. *Diabetologia*, **1982**, 22, 399-411.
- (3) Patel, M. P.; Patel, N. M.; Goyal, R. K. *Pharma Times*, **2006**, 38, 19-25.
- (4) Norberg, A.; Hoa, N. K.; Liepinsh, E.; Phan, D. V. Thuan, N. D.; Jornvall, H.; Sillard, R.; Ostenson, C.G. *J. Biol. Chem.* **2004**, 279, 41361-67.
- (5) Zemplén, G. *Ber. Dtsch Chem. Ges.*, **1926**, 59, 1254-1266.
- (6) Lipshutz, B. H.; Barton, J. C. *J. Org. Chem.* **1988**, 53, 4495-4499.
- (7) Madhusudan, S. K.; Agnihotri, G.; Negi, D. S.; Misra, A. K. *Carbohydr. Res.* **2005**, 340, 1373-1377.
- (8) Lewbart, M. L.; Schneider, J. J. *J. Org. Chem.* **1969**, 34, 3505-3512.
- (9) (a) Kaji, E.; Harita, N. *Tetrahedron Lett.* **2000**, 41, 53-56; (b) Kaji, E.; Shibayama, K.; In, K. *Tetrahedron Lett.* **2003**, 44, 4881-4885.
- (10) (a) Konradsson, P.; Udodong, U. E.; Fraser-Reid, B. *Tetrahedron Lett.* **1990**, 31, 4313-4316; (b) Veenemann, G. H.; van Leeuwen, S. H.; van Boom, J. H. *Tetrahedron Lett.* **1990**, 31, 1331-1334.
- (11) (a) Wang, W.; Kong, F. *Angew. Chem. Int. Ed. Engl.* **1999**, 38, 1247-1250; (b) Wang, W.; Kong, F. *J. Org. Chem.* **1998**, 63, 5744-5745.
- (12) Ziegler, T.; Echhardt, E.; Pantkowski, G. *J. Carbohydr. Chem.* **1994**, 13, 81-109.
- (13) Gurjar, M. K.; Adhikari, S. *Tetrahedron* **1997**, 53, 8629-8634.
- (14) Ferrier, R. J.; Furneaux, R. H. *J. Chem. Soc., Perkin Trans. 1*, **1977**, 1993-1996.
- (15) Pearlman, W. M. *Tetrahedron Lett.* **1967**, 8, 1663-1664.
- (16) Veeneman, G. H.; Gomes, L. J. F.; van Boom, J. H. *Tetrahedron* **1989**, 45, 7433-7448.



Scheme 1: Reagents: (a) 4-(OCH₃)-C₆H₅OH, TfOH, CH₂Cl₂, 2 h, 10 °C, 90 %; (b) (i) CH₃ONa, CH₃OH, r t, 3 h; (ii) 2,2-dimethoxypropane, *p*-toluenesulfonic acid, DMF, 6 h, 71% in two steps; (iii) benzyl bromide, tetrabutylammonium iodide, 50% aq. NaOH, CH₂Cl₂, r t, 3 h, 90%; (c) 80% aq. AcOH, 60 °C, 45 min, 93%; (d) (i) Bu₂SnO, CH₃OH, reflux, 6 h; (ii) benzoyl chloride, toluene, r t, 8 h, 80%; (e) *N*-iodosuccinimide, TfOH, CH₂Cl₂, -30 °C, 45 min, 78%; (f) 0.1 M CH₃ONa, CH₃OH, r t, 3 h, quantitative; (g) *N*-iodosuccinimide, TfOH, CH₂Cl₂, -30 °C, 45 min, 72%; (h) (i) CH₃ONa, CH₃OH, r t, 12 h; (ii) H₂, Pd(OH)₂-C, CH₃OH, r t, 24 h, 61% in two steps.



Scheme 2: Reagents: (a) *N*-Iodosuccinimide, TfOH, CH₂Cl₂, -30 °C, 45 min, 68%; (b) (i) CH₃ONa, CH₃OH, r t, 12 h; (ii) H₂, Pd(OH)₂-C, CH₃OH, r t, 10 h, 65% in two steps.